

# Isolated ovarian follicle culture: a promising strategy for fertility preservation

Sam Sudhakaran  
Roberto Gualtieri  
Vincenza Barbato  
Illaria Fiorentino  
Sabrina Braun  
Anna Merolla  
Riccardo Talevi

Department of Biology  
“Federico II” University, Napoli, Italy

## Address for correspondence:

Sam Sudhakaran, PhD  
Department of Biology  
“Federico II” University, Napoli, Italy  
E-mail: roschers007@gmail.com

## Summary

**Ovarian tissue cryopreservation represents one among the most preferred strategies for fertility preservation currently. However, concerns regarding the transmission of malignant cells during the transplantation of stored ovarian tissues, is a major restraint in recommending the procedure to patients diagnosed with all kinds of malignant disorders. On the contrary, use of isolated follicles for restoration of fertility in such patients could completely evade the possibility of cancer re-introduction after treatment. Follicles housed in the ovarian environment *in vivo* prevail under the mechanical and the chemical/nutritional support of the ovary. Although not complete, recent knowledge about the dynamics of follicular progression has led to improvements in the culture system adopted. This review aims at summarising the culture of isolated follicles *in vitro*, particularly emphasising the efforts made to mechanically and nutritionally support the follicle. Advances in follicular culture systems could**

**prove useful to highly improve the efficiency of current fertility restoration strategies and evade the concerns associated with the same.**

*KEY WORDS: fertility preservation, follicle culture, three dimensional support, isolated follicles.*

## Introduction

Cancer prevalence among women is till date, a major medical concern. Studies suggest that, 1 out of 51 women would have had an invasive cancer diagnosed by 39 years of age (1). Recent reports supported by the national cancer institute, have shown a 0.8% increase in cancer affected children during the past decade (2). However, recent advances in chemotherapy, radiotherapy, and bone marrow transplantation can cure as high as 90% of women and children affected by cancer and other disorders requiring such treatment. On the other side, ionizing radiation and aggressive chemotherapy can result in some degree of premature ovarian failure in almost 100% of patients requiring such therapy (3). Whole body irradiation coupled with intensive chemotherapy associated with bone marrow transplantation, poses one of the greatest threats to treated patients. The ovarian reserve is completely abolished after treatment regimes including alkylating agents such as busulphan (4). Moreover, several studies have shown that a radiation dose as low as 5- 20 Gy, is sufficient to cause gonadal function impairment (5-7).

## Concerns in routine ovarian tissue cryopreservation

Cryopreservation of the ovarian tissue is one of the mainstays in fertility preservation strategies adopted today among cancer affected women. The fact that ~25 live births have been reported till date using this procedure (8), could easily ar-

gue against considering this procedure under experimental phase any longer. Moreover, ovarian tissue cryopreservation is the only option available today for preservation of fertility in pre-pubertal girls and patients, requiring immediate initiation of potentially gonadotoxic anti-cancer treatment (9). Unfortunately, the most notable concern for ovarian tissue cryopreservation and transplant after the course of treatment, is the probability of re-introducing malignant cells back into the cured individual with very high chances of propagating the cancer again (10).

Ovarian metastases have been reported for most malignancies including breast cancer, lung cancer, renal tumors, neblastomas, Ewing's sarcoma, Hodgkins's lymphoma, Non-Hodgkin's lymphoma, biliary duct cancer and other gastrointestinal cancers (11-16). Studies evaluating the incidence of ovarian metastasis in different cancers show that ovarian involvement is highest in gastric cancer (55.8%), colon cancer (26.6%), breast cancer (24.2%), pulmonary carcinoma (23.4%), lymphoma (13.3%) uterine cancer (13.1%) and leukaemia (8.4%) (17).

Routine histological examination of the ovarian tissue fragments have proven to be inefficient in predicting the prevalence of malignant cells in the transplanted tissue (18). In this context, transplantation of stored ovarian tissue is particularly warranted in leukaemia, being a systemic disease with very high chances of metastasizing to the ovaries. In a study involving 18 leukaemia patients, routine histology and immunohistochemical analysis showed no presence of malignant cells in the biopsied ovarian tissue. However, highly sensitive reverse transcriptase PCR (RT-PCR) revealed the presence of molecular leukemic markers in the tissues of 9 out of 16 of these patients, previously thought to be safe from ovarian metastasis (18). Although, there is no conclusive report demonstrating the re-introduction of malignant cells through ovarian tissue transplanted after storage, studies in animal models have shown growth of intraperitoneal masses after transplantation of ovarian tissue from leukemic patients (18). Hence, transplantation of stored ovarian tissue in general can only be offered with extreme caution for re-restoration of fertility in cancer treated women. Considering these issues recent research in fertility preservation is focused on alternative approaches to circumvent this problem. Cryopreservation, culture and methods to re-implant isolated

ovarian follicles rather than whole tissue, is one among the most focussed topics in this context. Depending on age, the ovarian cortex houses thousands of dormant primordial follicles that can be isolated (19). These immature follicles represent the largest population of ovarian follicles and are more resistant to cryopreservation than advanced stage follicles (20). Furthermore, early stage follicles have shown to maintain normal morphology and ultrastructure following freezing, making them excellent candidates for long term preservation. The most important advantage of isolated follicle culture and transplant arises from the fact that the malignancy cannot cross the basal lamina of the follicle and the oocyte is protected from cancer cell invasion (21). Moreover, improvements in follicular development *in vitro* can also help these patients to obtain a larger number of oocytes for *in vitro* fertilization techniques, hence overcoming the concerns of re-transplantation all together. Henceforth, improvements in the isolation, culture and re-introduction of ovarian follicles can greatly improve current fertility preservation strategies and can provide a risk free method for the restoration of fertility in women affected by chemo/radiotherapy induced premature ovarian failure.

### Strategies for isolation of ovarian follicles

Several methods have been tried over the years for the successful isolation of ovarian follicles. Mechanical isolation of follicles have been in practice since the last 2 decades employing several instruments. The most commonly used techniques involve the use of tissue choppers (22), homogenizers (23), cell dissociation sieves (24) as reviewed by Valdevane et al. (25). Larger pre antral follicles have been isolated by microdissection using insulin needles (26). Mechanical isolation using fine needles has the advantage of preserving the basal lamina and the thecal layers intact maintaining the integrity of the follicle (27). A recent study successfully attained isolation of bovine primary follicles using mechanical disruption of ovarian cortex using a pasteur pipette. Furthermore, these follicles were demonstrated to survive *in vitro* culture for 21 days and form visible antral cavities (28). Enzymatic digestion has been the preferred method

of choice for the isolation of primordial/ primary follicles since these small follicles ranging (30-50um) are manually very hard to isolate. Although, enzymatic isolation can improve follicle yields compared to mechanical methods (29, 30), these aggressive enzymes can compromise follicular survival (27, 31, 32). A recent study observed spontaneous degeneration of isolated human follicles in culture, after enzymatic isolation. In spite of using a purified enzyme blend in place of traditional crude mixtures of collagenase, human follicular isolation could not be efficiently established using this enzymatic method (33). Furthermore, enzymatically isolated follicles particularly tend to lose their basal lamina causing granulosa cells to migrate away from the oocyte (34). Since isolation of follicles for culture is a crucial step in individual follicle culture, fine tuning the method and optimising the stage of follicle to be isolated is necessary for further development of this technique.

### **Towards a 3 dimensional culture of follicles *in vitro***

The most promising results in follicular development *in vitro* was attained by Eppig and O'Brien et al. in mouse model, producing live offspring from primordial follicles (35, 36). In spite of a decade of research following this initial success, similar results are yet to be obtained in larger mammals and human. Conventional 2 dimensional cultures fail to mimic the *in vivo* follicular environment. Culture of follicles directly on treated membranes or tissue culture surfaces destroys the spatial arrangement of follicles. Granulosa cells are seen to attach on the culture surface and cause follicular flattening (35). Growth of the oocyte and meiotic maturation relies on the signals exchange through gap junctions between the oocyte and surrounding granulosa cells (37). Loss of these gap junctions causes premature ovulation and degeneration of the oocyte (38). These signals are crucial for sharing paracrine factors that in turn promote growth of both cell types (39). This is supported by the fact that, oocyte is not able to transport amino acids and carry out glucose and cholesterol biosynthesis independently, in the absence of granulosa cell secreted factors (40).

### **Mimicing the *in vivo* follicular environment**

Recent efforts on improving the *in vitro* maintenance and growth of follicles can roughly said to be aimed at enhancing two most crucial factors that determine fate of the follicle *in vitro*.

#### **1) Physical/mechanical support to the growing follicle**

Research conducted in the murine model has demonstrated the importance of biomechanical environment in determining follicular growth and considers this physical support to be as crucial as the hormonal milieu of the follicle (41). To this end an array of different strategies has been adopted till date to spatially support the follicle and provide the physical stimuli that it needs to survive *in vitro* reviewed by (27). Use of V shaped microwell plates has seen to be useful to some extent in maintaining three dimensional architecture of bovine (42) and human follicles (43). Collagen owing to its natural presence in the extra cellular matrix surrounding the follicle has been used to support *in vitro* follicular growth since the emergence of 3 dimensional culture systems. Embedded culture of follicles in collagen (44) was found to be superior to collagen membrane inserts (45) in supporting follicular growth and architecture *in vitro*. Combelles et al. (46) demonstrated that follicles embedded in collagen matrix maintained their three dimensional architecture and demonstrated neuronal like extensions arising from the granulosa cells towards the oocyte. Recent success with the use of collagen in supporting antral cavity formation of early bovine primary follicles (28) indicates that collagen gels still have scope for research, to be considered an efficient compound to mechanically support *in vitro* follicle growth. However, shrinkage of the gel and decreased microscopic visibility over time are the most common problems faced with the use of collagen (29). Furthermore, the need for enzymatic treatment to dissolve the gel at the end of culture is another matter of concern (47). The most widely used three dimensional follicle encapsulation system till date is alginate (48, 49). Alginate encapsulation coupled with the use




Figure 1 - A) Isolated primordial follicles stained with Hoechst 33324 (nuclear stain): oocyte nucleus clearly visualized. B) Isolated secondary follicles stained with Hoechst 33324 (nuclear stain): oocyte nucleus clearly visualized.

of V shaped microwell plates have been tried to support human and (49) primate (50, 51) follicles. Encapsulation with alginate has the advantage of promoting nutrient, oxygen, hormone and growth factor exchange between the follicle and the culture medium. Although, the comparatively rigid nature of alginate has seen to inhibit follicular growth in mouse follicles (52), human (53) and primate follicles (54) seem to prefer the rigid support provided by the alginate matrix. Primate follicles have been successfully cultured over long periods in alginate gels (55). This could be directly accounted to the highly fibrous nature of the ovarian cortex seen in human, bovine and primate ovaries compared to mice. Incorporation of extracellular matrix components to alginate have been tried to develop synthetic matrices that improve the performance of alginate supported cultures (56). Studies on

the functions of extracellular matrix have revealed that it plays a crucial role in coordinating cell behaviour, cellular differentiation and secretion which are inevitable for follicular advancement (57). Our group demonstrated that alginate incorporated with collagen IV, a major extracellular component in ovary enhances the growth of isolated human follicles in culture. Moreover, ultrastructural analysis of these follicles revealed that culture of follicles embedded in alginate + collagen IV better preserved their three dimensional follicular architecture (58). Matrigel is a commercially available extracellular matrix composed of collagen IV, laminin, fibronectin, entactin, heparin sulphate and proteoglycans along with an array of growth factors (EGF, FGF, IGF-1, PDGF and TGF- $\beta$ ) that has been tried for the growth of follicles (53, 59, 60). The incorporation of bio-engineering has

led to the development of novel dynamic matrices in follicle culture. These compounds differ from traditional materials by their ability to more efficiently accommodate the drastically expanding follicle. Fibrin containing matrices are widely being tested recently, particularly after the recent success of developing a mature oocyte from a macaque primary follicle for the first time (61). Biodegradable matrices composed of Interpenetrating Networks of fibrin (IPN) (62) that intercalate with matrix proteins could better ensure sustained delivery of growth factors to the follicle (33). VEGF containing IPN matrix has recently been shown to support *in vitro* follicle growth, oocyte maturation and the subsequent development of a live offspring. Novel hydrogels comprising intercalating peptides that lyse in response to follicular proteases could dynamically allow expansion of the growing follicle. Hence, the future of designing matrices for the *in vitro* growth of follicles lies in incorporating concepts of bio-engineering and tissue culture to develop dynamic culture systems that adjusts to the changing dimensions of the follicle.

## 2) Chemical/nutritional demands of the follicle

A recent research has revealed that in order to extrapolate the success obtained in culturing rodent follicles into larger species a multi-step dynamic culture system is required to cater to the various transitional stages of mammalian follicular development (43, 63, 64). To this end, follicular development will have to address 3 major events: 1) Primordial follicle activation and initiation of growth, 2) Pre-antral to antral follicle transition, 3) Development of fertilizable oocytes from tertiary follicles (27).

Hence the primary step to be considered is the *in vitro* activation of primordial follicles. Unfortunately, there are no conclusive reports on the factors that control early follicle recruitment and growth. However, it seems safe to assume that the process should involve a complex interplay of inhibitory, stimulatory and maintenance factors (65). Recent research suggests a role of Phosphatidylinositol- 3-Kinase (PI3K) – Akt signalling pathway of the oocyte in kick starting the follicular growth (66). Fine tuning of the hormonal and chemical milieu of the follicle is

crucial to attain developmentally competent oocytes that complete cytoplasmic and nuclear maturation at a desired pace. Consequently several hormones and signalling molecules have been proposed to obtain optimal follicle growth *in vitro*. A few bio active components trialled have been discussed below.

Follicle Stimulating Hormone (FSH) is one among the most repeatedly trialled hormone to this end. FSH and estradiol are shown to have positive effects on antrum formation in granulosa cells in rodents (67) and pigs (68) for a long time. FSH along with LH activates cAMP system and activates the enzymes responsible for steroidogenesis in granulosa cells (69). FSH has also been shown to have positive effects on long term culture of bovine ovarian cortex in addition with GDF-9 and bFGF (70). Other studies have also reported the role of FSH in ensuring pre antral follicle survival and growth of primate (50) and human (53) follicles in 3 dimensional culture.

A bi-phasic requirement of FSH was revealed in a recent study that attained for the first time a 2 cell embryo from primate follicles cultured *in vitro* (51). Here they exposed follicles to high FSH levels before antrum formation and marginal levels after appearance of the antrum (51). In fact exposure to elevated levels of FSH for prolonged periods could disrupt the control of the oocyte over granulosa cell proliferation and differentiation (51) and lead to the pre-mature loss of trans zonal projections between oocytes and granulosa cells (71).

Epidermal Growth Factor (EGF) is yet another factor promoting pre antral follicle growth (72) as it induces granulosa cell proliferation and folliculogenesis (73) and progesterone synthesis by activating FSH receptors in granulosa cells. Positive effect of EGF has also been demonstrated in various animal models like pig (74), cow (75) and hamster (76).

Presence of Basic Fibroblast Growth Factor (bFGF) bioactivity is demonstrated in granulosa cells (77) and early growing follicles (78, 79). A combination of FSH, EGF and bFGF have recently gained acceptance as optimal media components that have given interesting results in the bovine model (28).

Activin protein has been localized in granulosa cells of human follicles. Hence, possible roles of activin as a follicular growth have been investigated (43, 71) based on the stimulatory effect of

activin on ovine pre-antral (80) and caprine follicles (81). This group demonstrated that activin not only does stimulate the growth of follicles but may also have functions in aiding the directional proliferation of granulosa cells by promoting the polarized expression of connexin proteins. Since loss of cellular polarization is a major threat *in vitro*, causing granulosa cells to proliferate undirectionally, activin could serve as an important modulator of cellular progression in the follicles. Activin was also seen to improve granulosa zona focal adhesions, the loss of which is one among the major concerns of *in vitro* follicular culture.

Fetuin, a glycoprotein component in serum and follicular fluid (82, 83) has been used to substitute the use of serum avoiding the concerns of using contaminated animal derivatives in culture. Schroeder et al. showed that fetuin increased zona pellucida solubility during oocyte maturation *in vitro* and supported a serum free culture environment (84). A protease inhibitory effect of fetuin was put forward for this observation. Fetuin has been suggested to improve cellular differentiation, growth and attachment *in vitro* (85, 86). Furthermore, fetuin was seen to maintain the integrity of alginate gels in long term culture (51).

Apart from the culture components oxygen tension is a key factor in determining the behaviour of any culture environment. Follicle culture *in vitro* has been regularly conducted at atmospheric oxygen tension (20% v/v /140 mm Hg) (50). Theoretically, follicles should be maintained at around 5% oxygen tension owing to the low partial pressure of oxygen in the peritoneal cavity i.e. 40mmHg (87). Low oxygen culture has been beneficial for the culture of rat pre antral follicles improving oocyte, viability maturation, parthenogenic activation and fertilization rates *in vitro* (88). Caprine pre natural follicles exhibited higher percentage of antrum formation at 5% oxygen as compared to 20% (89). Higher levels of reactive oxygen species are frequently associated with high partial pressure of oxygen and this oxidative stress induces cytotoxicity (90). Subsequently, culturing under low oxygen tension has seen to reduce cumulus cell apoptosis in canine oocyte cumulus complexes in culture (91). Lately, a higher number of healthy oocytes were also derived under low oxygen conditions, during *in vitro* culture of follicles (51).

## Conclusions

In spite of several years of research, many key factors determining the complex process of follicular maturation still remains a mystery. Further knowledge of these fundamental mechanisms are necessary to extrapolate these factors *in vitro* in turn making *in vitro* follicle maturation possible. Analysis on studies conducted till date reveals a thin balance of several signalling molecules and factors that fine tune the optimal growth of the follicular unit. Hence, studies aimed to optimize each progressive step in follicular maturation, to more closely mimic what occurs *in vivo*, would be needed to attain a successful *in vitro* follicular growth. Improvements made in culture conditions of follicles *in vitro* could eventually avoid the need to transplant whole ovarian tissues to patients opting for fertility preservation. Furthermore, ability to transplant individual follicles grown *in vitro* could eliminate concerns like ischemic damage, graft death and accelerated proliferation of follicles following transplant that exhausts the whole tissue in a single attempt. Hence, single follicle culture and transplant could aid in attaining numerous competent oocytes minimising the wastage of follicles, ensuring long term results for the patient.

## References

1. Altekruse SF, Kosary CL, Krapcho M, Neyman N, Aminou R, Waldron W. SEER cancer statistics review, 1975-2007. Bethesda, MD: National Cancer Institute.
2. Edwards K, Noone AM, Mariotto AB, Simard EP, Boscoe FP, Henley SJ, Jemal A, Cho H, Anderson RN, Kohler BA, Ehemann CR, Ward EM. Annual Report to the Nation on the Status of Cancer, 1975-2010, Featuring Prevalence of Comorbidity and Impact on Survival Among Persons With Lung, Colorectal, Breast, or Prostate Cancer, 2013.
3. Donnez J, Dolmans MM, Pellicer A, Diaz-Garcia C, Sanchez SM, Schmidt KT, Ernst E, Luyckx V, Andersen CY. Restoration of ovarian activity and pregnancy after transplantation of cryopreserved ovarian tissue: a review of 60 cases of reimplantation. *Fertil Steril.* 2013;99:1503-13.
4. Teinturier C, Hartmann O, Valteau-Couanet D, Benhamou E, Bougneres PF. Ovarian function after autologous bone marrow transplantation in childhood: high-dose busulfan is a major cause of ovarian failure. *Bone Marrow Transpl.* 1998;22:989-94.
5. Wallace WH, Thomson AB, Saran F, Kelsey TW. Predicting age of ovarian failure after radiation to a field that includes the ovaries. *Int J Radiat Oncol Biol Phys.*

- 2005;62:738-44.
6. Schmidt KT, Rosendahl M, Ernst E, Loft A, Andersen AN, Dueholm M. Autotransplantation of cryopreserved ovarian tissue in 12 women with chemotherapy-induced premature ovarian failure: the Danish experience. *Fertil Steril*. 2011;95:695-701.
  7. Anderson RA, Wallace WHB. Antimullerian hormone, the assessment of the ovarian reserve and the reproductive outcome of the young patient with cancer. *Fertil Steril*. 2013;99:1469-75.
  8. Campbell BK, Hernandez MJ, Onions V, Pincott AC, Aljaser F, Fisher J, McNeilly AS, Webb R, Picton HM. Restoration of ovarian function and natural fertility following the cryopreservation and autotransplantation of whole adult sheep ovaries. *Hum Reprod*. 2014;29:1749-63.
  9. Donnez J, Martinez MB, Jadoul P, Langendonck VA, Demylle D, Dolmans MM. Ovarian tissue cryopreservation and transplantation: a review. *Hum Reprod Update*. 2006;12:519-35.
  10. Chung K, Donnez J, Ginsburg E, Meirow D. Emergency IVF versus ovarian tissue cryopreservation: decision making in fertility preservation for female cancer patients. *Fertil Steril*. 2013;99:1534-42.
  11. Horvath TJ, Schindler AE. Ovarian metastases in breast carcinoma. *Fortschr Med*. 1977;95:358-60.
  12. Insabato L, De RG, Franco R, D'Onofrio V, Di VD. Ovarian metastasis from renal cell carcinoma: a report of three cases. *Int J Surg Pathol*. 2003;11:309-12.
  13. Khan MA, Dahill SW, Stewart KS. Primary Hodgkin's disease of the ovary. Case report. *Br J Obstet Gynaecol*. 1986;93:1300-1.
  14. Liu LY. Metastatic cancer in the ovary-report of 57 cases. *Zhonghua Zhong Liu Za Zhi*. 1989;11:464-7.
  15. Young RH, Kozakewich HP, Scully RE. Metastatic ovarian tumors in children: a report of 14 cases and review of the literature. *Int J Gynecol Pathol*. 1993;12:8-19.
  16. Hashimoto YN, Yamamoto T, Kamiura S, Seino H, Ohira H, Sawai K. Metastatic ovarian tumors: a review of 64 cases. *Gynecol Oncol*. 2003;89:314-7.
  17. Kyono K, Doshida M, Toya M, Sato Y, Akahira J, Sasano H. Potential indications for ovarian autotransplantation based on the analysis of 5,571 autopsy findings of females under the age of 40 in Japan. *Fertil Steril*. 2010;93:2429-30.
  18. Dolmans MM, Marinescu C, Saussoy P, Langendonck VA, Amorim C, Donnez J. Reimplantation of cryopreserved ovarian tissue from patients with acute lymphoblastic leukemia is potentially unsafe. *Blood* 2010;116:2908-14.
  19. Kristensen SG, Rasmussen A, Byskov AG, Andersen CY. Isolation of pre-antral follicles from human ovarian medulla tissue. *Hum Reprod*. 2011;26:157-66.
  20. Gougeon A. Dynamics of follicular growth in the human: a model from preliminary results. *Hum Reprod*. 1986;1:81-7.
  21. Abir R, Nitke S, Haroush AB, Fisch B. In vitro maturation of human primordial ovarian follicles: Clinical significance, progress in mammals, and methods for growth evaluation. *Histol Histopathol*. 2006;21:887-98.
  22. Figueiredo JR, Hulshof SC, Van den Hurk VR, Ectors FJ, Fontes RS, Nussgens B, Bevers MM, Beckers JF. Development of a combined new mechanical and enzymatic method for the isolation of intact preantral follicles from fetal, calf and adult bovine ovaries. *Theriogenology*. 1993;40:789-99.
  23. Nuttinck F, Mermillod P, Massip A, Dessy F. Characterization of in vitro growth of bovine preantral ovarian follicles: a preliminary study. *Theriogenology*. 1993;39:811-21.
  24. Jewgenow K, Goritz F. The recovery of preantral follicles from ovaries of domestic cats and their characterization before and after culture. *Anim Reprod Sci*. 1995;39:285-97.
  25. Valdevane RA, Melba OG, Figueiredo JR, Eduardo LG. In vitro culture of bovine preantral follicles: a review. *Reprod Biol Endocrinol*. 2014;12:78.
  26. Van den Hurk R, Spek ER, Hage WJ, Fair T, Ralph JH, Schotanus K. Ultrastructure and viability of isolated bovine preantral follicles. *Hum Reprod Update*. 1998;4:833-41.
  27. Telfer E, Zelinski MB. Ovarian follicle culture: advances and challenges for human and nonhuman primates. *Fertil Steril*. 2013;99:1534-42.
  28. Sun J, Li X. Growth and antrum formation of bovine primary follicles in long-term culture in vitro. *Reprod Biol*. 2013;13:221-8.
  29. Telfer E. The development of methods for isolation and culture of preantral follicles from bovine and porcine ovaries. *Theriogenology*. 1996;45:101-10.
  30. Hornick JE, Duncan FE, Shea LD, Woodruff TK. Multiple follicle culture supports primary follicle growth through paracrine-acting signals. *Reproduction*. 2013;145:19-32.
  31. Abir R, Roizman P, Fisch B, Nitke S, Okon E, Orvieto R, Rafael ZB. Pilot study of isolated early human follicles cultured in collagen gels for 24 hours. *Hum Reprod*. 1999;14:1299-301.
  32. Abir R, Fisch B, Nitke S, Okon E, Raz A, Rafael ZB. Morphological study of fully and partially isolated early human follicles. *Fertil Steril*. 2001;75:141-6.
  33. Laronda MM, Duncan FE, Hornick JE, Xu M, Pahnke JE, Whelan KA, Shea LD, Woodruff TK. Alginate encapsulation supports the growth and differentiation of human primordial follicles within ovarian cortical tissue. *J Assist Reprod Genet*. 2014;31:1013-28.
  34. Desai N, Alex A, AbdelHafez F, Calabro A, Goldfarb J, Fleischman A, Falcone T. Three-dimensional in vitro follicle growth: overview of culture models, biomaterials, design parameters and future directions. *Reprod. Biol Endocrinol*. 2010;8:119.
  35. Eppig JJ, O'Brien MJ. Development in vitro of mouse oocytes from primordial follicles. *Biol Reprod*. 1996;54:197-207.
  36. O'Brien MJ, Pendola JK, Eppig JJ. A revised protocol for in vitro development of mouse oocytes from primordial follicles dramatically improves their developmental competence. *Biol Reprod*. 2003;68:1682-6.
  37. Carabatsos MJ, Sellitto C, Goodenough DA, Albertini DF. Oocyte-granulosa cell heterologous gap junctions are required for the coordination of nuclear and cytoplasmic meiotic competence. *Dev Biol*. 2000;226:167-79.
  38. Eppig JJ, Pendola FL, Wigglesworth K, Pendola JK. Mouse oocytes regulate metabolic cooperativity between granulosa cells and oocytes: amino acid transport. *Biol Reprod*. 2005;73:351-7.
  39. Diaz FJ, Wigglesworth K, Eppig JJ. Oocytes are required for the preantral granulosa cell to cumulus cell transition in mice. *Dev Biol*. 2007;305:300-11.

40. Eppig JJ. Intercommunication between mammalian oocytes and companion somatic cells. *Bioessays*. 1991;13:569-74.
41. West ER, Xu M, Woodruff TK, Shea LD. Physical properties of alginate hydrogels and their effects on in vitro follicle development. *Biomaterials*. 2007;28:4439-48.
42. Thomas FH, Armstrong DG, Campbell BK, Telfer EE. Effects of insulin-like growth factor-1 bioavailability on bovine preantral follicular development in vitro. *Reproduction*. 2007;133:1121-8.
43. Telfer EE, McLaughlin M, Ding C, Thong KJ. A two step serum free culture system supports development of human oocytes from primordial follicles in the presence of activin. *Hum Reprod*. 2008;23:1151-8.
44. Loret de Mola JR, Barnhart K, Kopf GS, Heyner S, Gar-side W, Coutifaris CB. Comparison of two culture systems for the in-vitro growth and maturation of mouse preantral follicles. *Clin Exp Obstet Gynecol*. 2004;31:15-9.
45. Eppig JJ, Telfer EE. Isolation and culture of oocytes. *Methods Enzymol*. 1993;225:77-84.
46. Combelles CM, Fissore RA, Albertini DF, Racowsky C. In vitro maturation of human oocytes and cumulus cells using a co-culture three-dimensional collagen gel system. *Hum Reprod*. 2005;20:1349-58.
47. Torrance C, Telfer E, Gosden RG. Quantitative study of the development of isolated mouse pre-antral follicles in collagen gel culture. *J Reprod Fertil*. 1989;87:367-74.
48. Amorim CA, Van Langendonck A, David A, Dolmans MM, Donnez J. Survival of human pre-antral follicles after cryopreservation of ovarian tissue, follicular isolation and in vitro culture in a calcium alginate matrix. *Hum Reprod*. 2009;24:92-9.
49. Xu M, Banc A, Woodruff TK, Shea LD. Secondary follicle growth and oocyte maturation by culture in alginate hydrogel following cryopreservation of the ovary or individual follicles. *Biotechnol Bioeng*. 2009;103:378-86.
50. Xu J, Bernuci MP, Lawson MS, Yeoman RR, Fisher TE, Zelinski MB, Stouffer RL. Survival, growth, and maturation of secondary follicles from prepubertal, young and older adult, rhesus monkeys during encapsulated three-dimensional (3D) culture: effects of gonadotropins and insulin. *Reproduction*. 2010;140:685-97.
51. Xu J, Lawson MS, Yeoman RR, Zelinski MB, Stouffer RL. Secondary follicle growth and oocyte maturation during encapsulated three-dimensional culture in rhesus monkeys: effects of gonadotropins, oxygen and fetuin. *Hum Reprod*. 2011;26:1061-72.
52. Heise M, Koepsel R, Russell AJ, McGee EA. Calcium alginate microencapsulation of ovarian follicles impacts FSH delivery and follicle morphology. *Reprod Biol Endocrinol*. 2005;3:47.
53. Xu M, Barrett SL, West-Farrell E, Kondapalli LA, Kiewetter SE, Shea LD, Woodruff TK. In vitro grown human ovarian follicles from cancer patients support oocyte growth. *Hum Reprod*. 2009;24:2531-40.
54. Hornick JE, Duncan FE, Shea LD, Woodruff TK. Isolated primate primordial follicles require a rigid physical environment to survive and grow in vitro. *Hum Reprod*. 2012;27:1801-10.
55. Xu M, West-Farrell ER, Stouffer RL, Shea LD, Woodruff TK, Zelinski MB. Encapsulated three-dimensional culture supports development of nonhuman primate secondary follicles. *Biol Reprod*. 2009;81:587-94.
56. Kreeger PK, Woodruff TK, Shea LD. Murine granulosa cell morphology and function are regulated by a synthetic Arg-Gly-Asp matrix. *Mol Cell Endocrinol*. 2003;205:1-10.
57. Berkholtz CB, Shea LD, Woodruff TK. Extracellular matrix functions in follicle maturation. *Semin Reprod Med*. 2006;24:262-9.
58. Talevi R, Barbato V, Mollo V, Stefano DC, Finelli F, Ferraro R, Gualtieri R. Three dimensional alginate-collagen IV matrix enhances the in vitro growth of human isolated follicles. Abstracts of the 26th Annual Meeting of the European Society of Human Reproduction and Embryology, Rome, Italy. *Hum Reprod*. 2010;25(suppl 1):i260-77.
59. Hovatta O, Wright C, Krausz T, Hardy K, Winston RM. Human primordial, primary and secondary ovarian follicles in long-term culture: effect of partial isolation. *Hum Reprod*. 1999;14:2519-24.
60. Buyuk E. In vitro growth of mouse primary and early preantral follicles in a 3-dimensional culture system. *Fertil Steril*. 2003;80:79.
61. Xu J, Lawson MS, Yeoman RR, Molskness TA, Ting AY, Stouffer RL, Zelinski MB. Fibrin promotes development and function of macaque primary follicles during encapsulated three-dimensional culture. *Hum Reprod*. 2013;28:2187-200.
62. Shikanov A, Xu M, Woodruff TK, Shea LD. Interpenetrating fibrin-alginate matrices for in vitro ovarian follicle development. *Biomaterials*. 2009;30:5476-85.
63. Telfer EE, McLaughlin M. Human follicle activation and development in vitro. *Semin Reprod Med*. 2011;29:15-23.
64. Smits J, Dolmans MM, Donnez J, Fortune JE, Hovatta O, Jewgenow K. Current achievements and future research directions in ovarian tissue culture, in vitro follicle development and transplantation: implications for fertility preservation. *Hum Reprod Update*. 2010;16:395-414.
65. Nelson SM, Telfer EE, Anderson RA. The ageing ovary and uterus: new biological insights. *Hum Reprod Update*. 2013;19:67-83.
66. Reddy P, Liu L, Adhikari D, Jagarlamudi K, Rajareddy S, Shen Y. Oocyte specific deletion of Pten causes premature activation of the primordial follicle pool. *Science*. 2008;319:611-3.
67. Gore-Langton RE, Daniel SA. Follicle-stimulating hormone and estradiol regulate antrum-like reorganization of granulosa cells in rat preantral follicle cultures. *Biol Reprod*. 1990;43:65-72.
68. Hirao Y, Nagai T, Kubo M, Miyano T, Miyake M, Kato S. In vitro growth and maturation of pig oocytes. *J Reprod Fertil*. 1994;100:333-9.
69. Peng NN, Xue LQ, Ch KY. In vitro culture of follicular oocytes in mammals. *Journal of Hunan Agricultural University*. 1997;23:594-8.
70. Tang K, Yang WC, Li X, Wu CJ, Sang L, Yang LG. GDF-9 and bFGF enhance the effect of FSH on the survival, activation, and growth of cattle primordial follicles. *Anim Reprod Sci*. 2012;131:129-34.
71. McLaughlin M, Bromfield JJ, Albertini DF, Telfer EE. Activin promotes follicular integrity and oogenesis in cultured pre-antral bovine follicles. *Mol Hum Reprod*. 2010;6:644-53.
72. Celestino JJ, Bruno JB, Saraiva MV, Rocha RM, Brito

- IR, Duarte AB. Steady-state level of epidermal growth factor (EGF) mRNA and effect of EGF on in vitro culture of caprine preantral follicles. *Cell Tissue Res.* 2011;344:539-50.
73. Jewgenow K. Impact of peptide growth factors on the culture of small preantral follicles of domestic cats. *Theriogenology.* 1996;45:889-95.
74. Morbeck DE, Esbenshade KL, Flowers WL, Britt JH. Kinetics of follicle growth in the prepubertal gilt. *Biol Reprod.* 1992;47:485-91.
75. Wandji SA, Eppig JJ, Fortune JE. FSH and growth factors affect the growth and endocrine function in vitro of granulosa cells of bovine preantral follicles. *Theriogenology.* 1996;45:817-32.
76. Roy SK. Transforming growth factor-beta potentiation of follicle-stimulating hormone-induced deoxyribonucleic acid synthesis in hamster preantral follicles is mediated by a latent induction of epidermal growth factor. *Biol Reprod.* 1993;48:558-63.
77. Berisha B, Sinowatz F, Schams D. Expression and localization of fibroblast growth factor (FGF) family members during the final growth of bovine ovarian follicles. *Mol Reprod Dev.* 2004;67:162-71.
78. Wandji SA, Pelletier G, Sirard MA. Ontogeny and cellular localization of 125I-labeled basic fibroblast growth factor and 125I-labeled epidermal growth factor binding sites in ovaries from bovine fetuses and neonatal calves. *Biol Reprod.* 1992;47:807-13.
79. Wandji SA, Pelletier G, Sirard MA. Ontogeny and cellular localization of 125I-labeled insulin-like growth factor-I, 125I-labeled follicle-stimulating hormone, and 125I-labeled human chorionic gonadotropin binding sites in ovaries from bovine fetuses and neonatal calves. *Biol Reprod.* 1992;47:814-22.
80. Thomas FH, Armstrong DG, Telfer EE. Activin promotes oocyte development in ovine preantral follicles in vitro. *Reprod Biol Endocrinol.* 2003;1:76.
81. Silva JR, Tharasanit T, Taverne MA, van der Weijden GC, Santos RR, Figueiredo JR, van den Hurk R. The activin-follistatin system and in vitro early follicle development in goats. *J Endocrinol.* 2006;189:113-25.
82. Høyer PE, Terkelsen OB, Byskov GA, Nielsen H. Fetuin and fetuin messenger RNA in granulosa cells of the rat ovary. *Biol Reprod.* 2001;65:1655-62.
83. Kalab P, Shultz RM, Kopf GS. Modifications of the mouse zona pellucida during oocyte maturation: inhibitory effects of follicular fluid, fetuin, and  $\alpha$ 2HS-glycoprotein. *Biol Reprod.* 1993;49:561-7.
84. Schroeder AC, Schultz RM, Kopf GS, Taylor FR, Becker RB, Eppig JJ. Fetuin inhibits zona pellucida hardening, conversion of ZP2 to ZP2f during spontaneous mouse oocyte maturation in vitro in the absence of serum. *Biol Reprod.* 1990;43:891-7.
85. Nie Z. Fetuin. its enigmatic property of growth promotion. *Am J Physiol.* 1992;263:C551-62.
86. Demetriou M, Binkert C, Sukhu B, Tenenbaum HC, Dennis JW. Fetuin/  $\alpha$ 2-HS glycoprotein is a transforming growth factor- $\beta$  type II receptor mimic and cytokine antagonist. *J Biol Chem.* 1996;271:12755-61.
87. Tsai AG, Friesenecker B, Mazzoni MC, Kerger H, Buerk DG, Johnson PC, Intaglietta M. Microvascular and tissue oxygen gradients in the rat mesentery. *Proc Natl Acad Sci. USA* 1998;95:6590-5.
88. Heise MK, Koepsel R, McGee EA, Russell AJ. Dynamic oxygen enhances oocyte maturation in long-term follicle culture. *Tissue Eng Part C Methods.* 2009;15:323-32.
89. Silva CM, Matos MH, Rodrigues GQ, Faustino LR, Pinto LC, Chaves RN, Araujo VR, Campello CC, Figueiredo JR. In vitro survival and development of goat preantral follicles in two different oxygen tensions. *Anim Reprod Sci.* 2010;117:83-9.
90. Evans MD, Dizdaroglu M, Cooke MS. Oxidative DNA damage and disease: induction, repair and significance. *Mutat Res.* 2004;567:1-61.
91. Silva AE, Rodriguez P, Cavalcante LF, Rodrigues BA, Rodrigues JL. The influence of oxygen tension on cumulus cell viability of canine COCs matured in high-glucose medium. *Reprod Domest Anim.* 2009;44:259-62.